

## LOCAL ILL-POSEDNESS OF THE 1D ZAKHAROV SYSTEM

JUSTIN HOLMER

ABSTRACT. Ginibre-Tsutsumi-Velo (1997) proved local well-posedness for the Zakharov system

$$\begin{aligned} i\partial_t u + \Delta u &= nu \\ \partial_t^2 n - \Delta n &= \Delta |u|^2 \\ u(x, 0) &= u_0(x), \\ n(x, 0) &= n_0(x), \quad \partial_t n(x, 0) = n_1(x) \end{aligned}$$

where  $u = u(x, t) \in \mathbb{C}$ ,  $n = n(x, t) \in \mathbb{R}$ ,  $x \in \mathbb{R}$ , and  $t \in \mathbb{R}$ . The proof was made for any dimension  $d$ , in the inhomogeneous Sobolev spaces  $(u, n) \in H^k(\mathbb{R}^d) \times H^s(\mathbb{R}^d)$  for a range of exponents  $k, s$  depending on  $d$ . Here we restrict to dimension  $d = 1$  and present a few results establishing local ill-posedness for exponent pairs  $(k, s)$  outside of the well-posedness regime. The techniques employed are rooted in the work of Bourgain (1993), Birnir-Kenig-Ponce-Svanstedt-Vega (1996), and Christ-Colliander-Tao (2003) applied to the nonlinear Schrödinger equation.

### 1. INTRODUCTION

In this paper, we examine the one-dimensional Zakharov system (1DZS)

$$\begin{aligned} i\partial_t u + \partial_x^2 u &= nu \\ \partial_t^2 n - \partial_x^2 n &= \partial_x^2 |u|^2, \\ u(x, 0) &= u_0(x), \\ n(x, 0) &= n_0(x), \quad \partial_t n(x, 0) = n_1(x) \end{aligned} \tag{1.1}$$

where  $u = u(x, t) \in \mathbb{C}$ ,  $n = n(x, t) \in \mathbb{R}$ ,  $x \in \mathbb{R}$ , and  $t \in \mathbb{R}$ . Local well-posedness in the inhomogeneous Sobolev spaces  $(u, n) \in H^k(\mathbb{R}) \times H^s(\mathbb{R})$  has been obtained by means of the contraction method in the Bourgain space

$$\|u\|_{X_{k,b_1}^s} = \left( \iint_{\xi, \tau} \langle \xi \rangle^{2k} \langle \tau + |\xi|^2 \rangle^{2b_1} |\hat{u}(\xi, \tau)|^2 d\xi d\tau \right)^{1/2}$$

---

2000 *Mathematics Subject Classification*. 35Q55, 35Q51, 35R25.

*Key words and phrases*. Zakharov system; Cauchy problem; local well-posedness; local ill-posedness.

©2007 Texas State University - San Marcos.

Submitted March 24, 2006. Published February 12, 2007.

Partially supported by an NSF postdoctoral fellowship.

by Bourgain-Colliander [1] and Ginibre-Tsutsumi-Velo [7].<sup>1</sup> In the latter paper, the following result is obtained.

**Theorem 1.1** ([7, Prop. 1.2]). *Problem (1.1) is locally well-posed for initial data  $(u_0, n_0, n_1) \in H^k \times H^s \times H^{s-1}$  provided that*

$$\begin{aligned} k &\geq 0, & s &\geq -\frac{1}{2}; \\ -1 &\leq s - k < \frac{1}{2}, & s &\leq 2k - \frac{1}{2} \end{aligned}$$

*Specifically:*

- (1) Existence. *For all  $R > 0$ , if  $\|u_0\|_{H^k} + \|n_0\|_{H^s} + \|n_1\|_{H^{s-1}} < R$ , then there exist  $T = T(R)$  and a solution  $(u, n)$  to (1.1) on  $[0, T]$  such that*

$$\begin{aligned} \|u\|_{C([0, T]; H_x^k)} &\leq c\|u_0\|_{H^k}, \\ \|n\|_{C([0, T]; H_x^s)} + \|\partial_t n\|_{C([0, T]; H_x^{s-1})} &\leq c(\|u_0\|_{H^k})^2(\|n_0\|_{H^s} + \|n_1\|_{H^{s-1}}) \end{aligned}$$

*and  $u \in X_{k, b_1}^S$ , where  $b_1$  is given by Table 1.*

- (2) Uniqueness.<sup>2</sup> *This solution is unique among solutions  $(u, n)$  such that*

$$u \in C([0, T]; H_x^k) \cap X_{k, b_1}^S.$$

- (3) Uniform continuity of the data-to-solution map. *For a fixed  $R > 0$ , taking  $T = T(R)$  as above, the map  $(u_0, n_0, n_1) \mapsto (u, n, \partial_t n)$  as a map from the  $R$ -ball in  $H^k \times H^s \times H^{s-1}$  to  $C([0, T]; H_x^k) \times C([0, T]; H_x^s) \times C([0, T]; H_x^{s-1})$  is uniformly continuous.*

The region of local well-posedness in this theorem is depicted in Fig. 1. We shall outline the [7] proof of Theorem 1.1 in §2 since the estimates are needed in the proof of Theorem 1.2 in §3.

Our goal in this paper is to establish local ill-posedness outside of the well-posedness strip, in particular near the optimal corner  $k = 0$ ,  $s = -1/2$ . That is, we consider the region (1)  $s > 2k - \frac{1}{2}$  (above the strip), and (2)  $s < -1/2$  (below the strip). In the first region, the wave data  $(n_0, n_1)$  is somewhat smoother than the Schrödinger data  $u_0$ . As a result, the forcing term  $\partial_x^2 |u|^2$  of the wave equation, as time evolves, introduces disturbances that are rougher than the wave data, and the wave solution  $n$  does not retain its higher initial regularity. This is quantified in Theorem 1.2 below. In the second region, the Schrödinger data  $u_0$  is somewhat smoother than the wave data  $(n_0, n_1)$ . As a result, the forcing term  $nu$  of the Schrödinger equation introduces disturbances that are rougher than the Schrödinger data, and the Schrödinger solution  $u$  does not retain its higher initial regularity. This is quantified in Theorem 1.3 and 1.4 below. These simplistic explanations are, at least, accurate for  $k > 0$ . For  $k < 0$ , there are possibly multiple simultaneous causes for breakdown, although we find that our methods still yield information in this setting.

<sup>1</sup>Actually, these papers consider, more generally, the system in dimensions  $d = 2, 3$  and  $d \geq 1$ , respectively.

<sup>2</sup>(1.1) can be recast as an integral equation in  $\bar{u}$  alone with  $W(n_0, n_1)$  solving (2.2) appearing as a coefficient. Then,  $n$  can be expressed in terms of  $u$  and  $W(n_0, n_1)$ , and therefore  $n$  need not enter into the uniqueness claim.




FIGURE 1. The enclosed gray-shaded strip, which extends infinitely to the upper-right, gives the set of pairs  $(k, s)$  for which well-posedness has been established by [7] (see Theorem 1.1) for  $(u_0, n_0, n_1) \in H^k \times H^s \times H^{s-1}$ . Solid lines are included in the well-posedness region, while the dashed line is not. Theorem 1.2 provides an ill-posedness result of type “norm inflation in  $n$ ” inside the region bounded by the horizontal dotted line  $s = -1/2$ , the slanted line  $s = 2k - \frac{1}{2}$ , and the vertical dotted line  $k = 1$ . Theorem 1.3 provides an ill-posedness result of type “phase decoherence in  $u$ ” along the solid vertical line extending down from the point  $(0, -3/2)$ .

We will draw upon and suitably modify techniques developed by Birnir-Kenig-Ponce-Svanstedt-Vega [2], Christ-Colliander-Tao [5], and Bourgain [3], who addressed ill-posedness issues for the nonlinear Schrödinger equation. For a survey of ill-posedness results for nonlinear dispersive equations, see Tzvetkov [10].

Our first result demonstrates that the boundary line  $s \leq 2k - \frac{1}{2}$  in Theorem 1.1 is sharp.

**Theorem 1.2.** *Let  $0 < k < 1$  and  $s > 2k - \frac{1}{2}$  or  $k \leq 0$  and  $s > -1/2$ . There exists a sequence  $\phi_N \in \mathcal{S}$  such that  $\|\phi_N\|_{H^k} \leq 1$  for all  $N$  and the corresponding solution  $(u_N, n_N)$  to (1.1) on  $[0, T]$  with initial data  $(\phi_N, 0, 0)$  satisfies*

$$\|n_N(t)\|_{H_x^s} \geq ctN^\alpha \quad \text{for } 0 < t \leq T, \quad N \geq ct^{-1} \quad (1.2)$$

where  $\alpha = \alpha(k, s) > 0$ . The time interval  $[0, T]$  here is independent of  $N$ .

The form of ill-posedness appearing in Theorem 1.2 is referred to as “norm inflation”. The result is first reduced to the case where  $k > 0$  and  $s$  is just above the line  $s = 2k - \frac{1}{2}$ . In this case, Theorem 1.1 applied with  $s = 2k - \frac{1}{2}$  (the wave initial data is 0) provides the existence of a solution  $(u_N, n_N)$  on a time interval  $T$ , independent of  $N$ , with uniform-in- $N$  control on  $\|u_N\|_{X_{k,b_1}^s}$ . The estimates of [7] will enable us to show that  $u_N$  is comparable to  $e^{it\partial_x^2}\phi_N$  in a slightly stronger norm than  $X_{k,b_1}^s$  (on this fixed in  $N$  time interval) and then Theorem 1.2 follows from the fact that (1.2) holds with  $n_N = \square^{-1}\partial_x^2|u_N|^2$  replaced by  $\square^{-1}\partial_x^2|e^{it\partial_x^2}\phi_N|^2$ , which can be directly verified.<sup>3</sup> The proof is given in §3.

Our second theorem demonstrates lack of uniform continuity of the data-to-solution map, for any  $T > 0$ , as a map from the unit ball in  $H^k \times H^s \times H^{s-1}$  to  $C([0, T]; H^k) \times C([0, T]; H^s) \times C([0, T]; H^{s-1})$  for  $k = 0$  and any  $s < -\frac{3}{2}$ . We first show that if one issue is ignored, we can, in a manner similar to [2], make use of an explicit soliton class to demonstrate that for any  $T > 0$  there are two waves, close in amplitude on all of  $[0, T]$ , initially of the same phase but that slide completely out of phase by time  $T$ . This form of ill-posedness is termed “phase decoherence”. The soliton class for (1.1) that we use appears in [8] [11]. The “ignored issue” pertains to low frequencies of  $n_0(x)$ , and can be resolved by invoking the method of [5] to construct a “near soliton” class offering more flexibility than the exact explicit soliton class in the selection of  $n_0(x)$ . This is, however, not straightforward since (1.1) lacks scaling and Galilean invariance, which was used to manufacture the solution class in [5].

**Theorem 1.3.** *Suppose  $k = 0$ ,  $s < -\frac{3}{2}$ . Fix any  $T > 0$  and  $\delta > 0$ . Then there is a pair of Schwartz class initial data tuples  $(u_0, n_0, 0)$  and  $(\tilde{u}_0, \tilde{n}_0, 0)$  giving rise to solutions  $(u, n)$  and  $(\tilde{u}, \tilde{n})$  on  $[0, T]$  such that the data is of unit size*

$$\|u_0\|_{H^k}, \|n_0\|_{H^s} \sim 1, \quad \|\tilde{u}_0\|_{H^k}, \|\tilde{n}_0\|_{H^s} \sim 1$$

and initially close

$$\|u_0 - \tilde{u}_0\|_{H^k} + \|n_0 - \tilde{n}_0\|_{H^s} \leq \delta$$

but the solutions become well-separated by time  $T$  in the Schrödinger variable

$$\|u(\cdot, t) - \tilde{u}(\cdot, t)\|_{L_{[0,T]}^\infty H_x^k} \sim 1.$$

We expect that this result can be extended to all  $k \in \mathbb{R}$  and  $s < -3/2$ , although preliminary efforts were abandoned since the computations became very lengthy and technical. The proof of Theorem 1.3 appears in §5.

Our final theorem employs a method of Bourgain [3].

<sup>3</sup> $w = \square^{-1}f$  is the solution to  $\square w = (\partial_t^2 - \partial_x^2)w = f$ ,  $w(x, 0) = 0$ ,  $\partial_t w(x, 0) = 0$ .

**Theorem 1.4.** *For any  $T > 0$ , the data-to-solution map, as a map from the unit ball in  $H^k \times H^s \times H^{s-1}$  to  $C([0, T]; H^k) \times C([0, T]; H^s) \times C([0, T]; H^{s-1})$  fails to be  $C^2$  for  $k \in \mathbb{R}$  and  $s < -1/2$ .*

This is a weaker form of ill-posedness than the phase decoherence of Theorem 1.3, although it covers the full region below the well-posedness boundary  $s = -1/2$  of [7]. The proof is given in §6.

## 2. THE LOCAL THEORY

We outline and review the local well-posedness argument in [7] since the estimates will be needed in the proofs of Theorems 1.2, 1.4.

Let  $[U(t)u_0](\xi) = e^{-it\xi^2} \hat{u}_0(\xi)$  and

$$U *_R f(\cdot, t) = \int_0^t U(t-t')f(t') dt'$$

denote the Schrödinger group and Duhamel operators, respectively. Define the Schrödinger Bourgain spaces  $X_{k,\alpha}^S, Y_k^S$  by the norms

$$\begin{aligned} \|z\|_{X_{k,\alpha}^S} &= \left( \iint_{\xi,\tau} \langle \xi \rangle^{2k} \langle \tau + |\xi|^2 \rangle^{2\alpha} |\hat{z}(\xi, \tau)|^2 d\xi d\tau \right)^{1/2} \\ \|z\|_{Y_k^S} &= \left( \int_{\xi} \langle \xi \rangle^{2k} \left( \int_{\tau} \langle \tau + |\xi|^2 \rangle^{-1} |\hat{z}(\xi, \tau)| d\tau \right)^2 d\xi \right)^{1/2}. \end{aligned} \quad (2.1)$$

Consider an initial wave data pair  $(n_0, n_1)$ . Split  $n_1 = n_{1L} + n_{1H}$  into low and high frequencies<sup>4</sup>, and set  $\hat{\nu}(\xi) = \frac{\hat{n}_{1H}(\xi)}{i\xi}$ , so that  $\partial_x \nu = n_{1H}$ . Let

$$\begin{aligned} W_+(n_0, n_1)(x, t) &= \frac{1}{2}n_0(x-t) - \frac{1}{2}\nu(x-t) + \frac{1}{2} \int_{x-t}^x n_{1L}(y) dy \\ W_-(n_0, n_1)(x, t) &= \frac{1}{2}n_0(x+t) + \frac{1}{2}\nu(x+t) + \frac{1}{2} \int_x^{x+t} n_{1L}(y) dy \end{aligned}$$

so that

$$\begin{aligned} (\partial_t \pm \partial_x)W_{\pm}(n_0, n_1)(x, t) &= \frac{1}{2}n_{1L}(x) \\ W_{\pm}(n_0, n_1)(x, 0) &= \frac{1}{2}n_0(x) \mp \frac{1}{2}\nu(x). \end{aligned}$$

By setting  $n = W_+(n_0, n_1) + W_-(n_0, n_1)$ , we obtain a solution to the linear homogeneous problem

$$\begin{aligned} \partial_t^2 n - \partial_x^2 n &= 0 \quad t, x \in \mathbb{R} \\ n(x, 0) = n_0(x), \quad \partial_t n(x, 0) &= n_1(x) \quad n = n(t, x) \in \mathbb{R} \end{aligned} \quad (2.2)$$

Let

$$W_{\pm} *_R f(x, t) = \frac{1}{2} \int_0^t f(x \mp s, t-s) ds \quad (2.3)$$

so that

$$(\partial_t \pm \partial_x)W_{\pm} *_R f(x, t) = \frac{1}{2}f(x, t), \quad W_{\pm} f(x, 0) = 0, \quad \partial_t W_{\pm} f(x, 0) = \frac{1}{2}f(x, 0).$$

<sup>4</sup>This decomposition is needed, for otherwise the estimate in Lemma 2.1(2) would have to be modified to have  $\|n_1\|_{H^s}$  in place of  $\|n_1\|_{H^{s-1}}$  on the right-hand side

It follows that if we set  $n = W_- *_{\mathbb{R}} f - W_+ *_{\mathbb{R}} f$ , then we obtain a solution to the linear inhomogeneous problem

$$\begin{aligned} \partial_t^2 n - \partial_x^2 n &= \partial_x f \quad t, x \in \mathbb{R} \\ n(x, 0) = 0, \partial_t n(x, 0) &= 0 \quad n(x, t) \in \mathbb{R} \end{aligned}$$

Define the one-dimensional reduced wave Bourgain spaces  $X_{s,\alpha}^{W\pm}, Y_s^{W\pm}$  as

$$\begin{aligned} \|z\|_{X_{s,\alpha}^{W\pm}} &= \left( \iint_{\xi,\tau} \langle \xi \rangle^{2s} \langle \tau \pm \xi \rangle^{2\alpha} |\hat{z}(\xi, \tau)|^2 d\xi d\tau \right)^{1/2} \\ \|z\|_{Y_s^{W\pm}} &= \left( \int_{\xi} \langle \xi \rangle^{2s} \left( \int_{\tau} \langle \tau \pm \xi \rangle^{-1} |\hat{z}(\xi, \tau)| d\tau \right)^2 d\xi \right)^{1/2}. \end{aligned} \quad (2.4)$$

Let  $\psi(t) = 1$  on  $[-1, 1]$  and  $\psi(t) = 0$  outside of  $[-2, 2]$ . Let  $\psi_T(t) = \psi(t/T)$ , which will serve as a time cutoff for the Bourgain space estimates. For clarity, we write  $\psi_1(t) = \psi(t)$ . We can now recast (1.1) as

$$\begin{aligned} i\partial_t u + \partial_x^2 u &= (n_+ + n_-)u \quad x \in \mathbb{R}, t \in \mathbb{R} \\ (\partial_t \pm \partial_x)n_{\pm} &= \mp \frac{1}{2} \partial_x |u|^2 + \frac{1}{2} n_{1L} \end{aligned} \quad (2.5)$$

where  $n = n_+ + n_-$ , which has the integral equation formulation

$$\begin{aligned} u(t) &= U(t)u_0 - iU *_{\mathbb{R}} [(n_+ + n_-)u](t) \\ n_{\pm}(t) &= W_{\pm}(t)(n_0, n_1) \mp W_{\pm} *_{\mathbb{R}} (\partial_x |u|^2)(t). \end{aligned}$$

**Lemma 2.1** (Group estimates).

- (1) Schrödinger.  $\|\psi_1(t)U(t)u_0\|_{X_{k,b_1}^S} \lesssim \|u_0\|_{H^k}$ .
- (2) 1-d Wave.  $\|\psi_1(t)W_{\pm}(t)(n_0, n_1)\|_{X_{s,b}^{W\pm}} \lesssim \|n_0\|_{H_x^s} + \|n_1\|_{H_x^{s-1}}$ .

**Lemma 2.2** (Duhamel estimates). *Suppose  $T \leq 1$ .*

- (1) Schrödinger. *If  $0 \leq c_1 < \frac{1}{2}$ ,  $0 \leq b_1$ ,  $b_1 + c_1 \leq 1$ , then  $\|\psi_T U *_{\mathbb{R}} f\|_{X_{k,b_1}^S} \lesssim T^{1-b_1-c_1} \|f\|_{X_{k,-c_1}^S}$ .  
If  $0 \leq b_1 \leq \frac{1}{2}$ , then  $\|\psi_T U *_{\mathbb{R}} f\|_{X_{k,b_1}^S} \lesssim T^{\frac{1}{2}-b_1} (\|f\|_{X_{k,-\frac{1}{2}}^S \cap Y_k^S})$ .  
 $\|U *_{\mathbb{R}} f\|_{C(\mathbb{R}_t; H_x^k)} \lesssim \|f\|_{Y_k^S}$ .*
- (2) 1-d Wave. *If  $0 \leq c < \frac{1}{2}$ ,  $0 \leq b$ ,  $b + c \leq 1$ , then  $\|\psi_T W_{\pm} *_{\mathbb{R}} f\|_{X_{s,b}^{W\pm}} \lesssim T^{1-b-c} \|f\|_{X_{s,-c}^{W\pm}}$ .  
If  $0 \leq b \leq \frac{1}{2}$ , then  $\|\psi_T W_{\pm} *_{\mathbb{R}} f\|_{X_{s,b}^{W\pm}} \lesssim T^{\frac{1}{2}-b} (\|f\|_{X_{s,-\frac{1}{2}}^{W\pm} \cap Y_s^{W\pm}})$ .  
 $\|W_{\pm} *_{\mathbb{R}} f\|_{C(\mathbb{R}_t; H_x^s)} \lesssim \|f\|_{Y_s^{W\pm}}$ .*

**Lemma 2.3** ([7, Lemma 4.3/4.5]). *Let  $k, s, b, c_1, b_1$  satisfy*

$$\begin{aligned} s &\geq -\frac{1}{2}, \quad k \geq 0, \quad s - k \geq -1, \\ b, c_1, b_1 &> \frac{1}{4}, \quad b + c_1 > \frac{3}{4}, \quad b + b_1 > \frac{3}{4}, \\ s - k &\geq -2c_1 \end{aligned}$$

*Then*

$$\|n_{\pm} u\|_{X_{k,-c_1}^S \cap Y_k^S} \lesssim \|n_{\pm}\|_{X_{s,b}^{W\pm}} \|u\|_{X_{k,b_1}^S}.$$

$s - k = -1$	$b_1 = \frac{1}{2} - \epsilon$ $c_1 = \frac{1}{2}$	$b = \frac{3}{4} - 3\epsilon$ $c = \frac{1}{4} + 2\epsilon$
$-1 < s - k < -1/2$	$b_1 = \frac{s-k}{2} + 1 - \epsilon$ $c_1 = -\frac{s-k}{2}$	$b = \frac{3}{4} - 2\epsilon$ $c = \frac{1}{4} + \epsilon$
$-\frac{1}{2} \leq s - k \leq 0$	$b_1 = \frac{3}{4} - 2\epsilon$ $c_1 = \frac{1}{4} + \epsilon$	$b = \frac{3}{4} - 2\epsilon$ $c = \frac{1}{4} + \epsilon$
$0 \leq s - k < \frac{1}{2}$	$b_1 = \frac{3}{4} - 2\epsilon$ $c_1 = \frac{1}{4} + \epsilon$	$b = \frac{3}{4} - \frac{s-k}{2} - 2\epsilon$ $c = \frac{s-k}{2} + \frac{1}{4} + \epsilon$

TABLE 1. Values of  $b_1, c_1, b, c$  meeting the criteria of Lemmas 2.3, 2.4 for various intervals of  $s - k$ . Note that  $b_1 + c_1 \leq 1 - \epsilon$  and  $b + c \leq 1 - \epsilon$  in order to capture a factor  $T^\epsilon$  from Lemma 2.2. Also note that  $b_1, b > \frac{1}{2}$  and  $c_1, c < \frac{1}{2}$  for all cases except  $s - k = -1$ .

**Lemma 2.4** ([7, Lemma 4.4/4.6]). *Let  $k, s, c, b_1$  satisfy*

$$\begin{aligned} s - 2k &\leq -\frac{1}{2}, & k &\geq 0, & s - k &< \frac{1}{2}, \\ c, b_1 &> \frac{1}{4}, & c + b_1 &> \frac{3}{4}, \\ s - k &\leq 2b_1 - 1, & s - k &< 2c - \frac{1}{2} \end{aligned}$$

Then

$$\|\partial_x(u_1 \bar{u}_2)\|_{X_{s,-c}^{W_\pm} \cap Y_s^{W_\pm}} \lesssim \|u_1\|_{X_{k,b_1}^S} \|u_2\|_{X_{k,b_1}^S}.$$

To obtain Theorem 1.1, fix  $0 < T < 1$ , and consider the maps  $\Lambda_S, \Lambda_{W_\pm}$

$$\Lambda_S(u, n_\pm) = \psi_1 U u_0 + \psi_T U *_R [(n_+ + n_-)u] \quad (2.6)$$

$$\Lambda_{W_\pm}(u) = \psi_1 W_\pm(n_0, n_1) \pm \psi_T W_\pm *_R (\partial_x |u|^2). \quad (2.7)$$

For  $T = T(\|u_0\|_{H^k}, \|n_0\|_{H^s}, \|n_1\|_{H^{s-1}})$ , a fixed point

$$(u(t), n_\pm(t)) = (\Lambda_S(u, n_\pm), \Lambda_{W_\pm}(u))$$

is obtained in  $X_{k,b_1}^S \times X_{s,b}^{W_\pm}$  satisfying

$$\|u\|_{X_{k,b_1}^S} \lesssim \|u_0\|_{H^k} \quad (2.8)$$

$$\|n\|_{X_{s,b}^{W_\pm}} \lesssim \|n_0\|_{H^s} + \|n_1\|_{H^{s-1}} + \|u_0\|_{H^k}^2 \quad (2.9)$$

by applying Lemmas 2.1, 2.2, 2.3, 2.4 with values for  $b_1, c_1, b, c$  given by Table 1.

Consider first the case  $s - k > -1$ . We note from Table 1 that  $b_1, b > \frac{1}{2}$ , and thus we have the Sobolev imbeddings

$$\begin{aligned} \|u\|_{C(\mathbb{R}_t; H_x^k)} &\lesssim \|u\|_{X_{k,b_1}^S} \\ \|n_\pm\|_{C(\mathbb{R}_t; H_x^s)} &\lesssim \|n_\pm\|_{X_{s,b}^{W_\pm}}. \end{aligned} \quad (2.10)$$

Also,

$$\partial_t n(x, t) = \partial_t(n_+ + n_-)(x, t) = \partial_x(-n_+ + n_-)(x, t) + n_{1L}(x)$$

and thus

$$\|\partial_t n\|_{C(\mathbb{R}_t; H_x^{s-1})} \lesssim \|n_\pm\|_{X_{s,b}^{W_\pm}} + \|n_1\|_{H^{s-1}}. \quad (2.11)$$

Similar estimates apply to differences of solutions.

Consider now the case  $s - k = -1$ , where it is necessary to take  $b_1 < \frac{1}{2}$ . We return to (2.6) and estimate directly using Lemma 2.2 to obtain

$$\|u\|_{C(\mathbb{R}_t; H_x^k)} \lesssim \|u_0\|_{H^k} + \|n_{\pm}u\|_{Y_k^s}$$

and by Lemma 2.3,

$$\|n_{\pm}u\|_{Y_k^s} \lesssim \|n_{\pm}\|_{X_{k,b}^{w_{\pm}}} \|u\|_{X_{s,b_1}^s}$$

where  $b_1, b$  are as specified in the Table 1, and the right-hand side is appropriately bounded by (2.8), (2.9). The bounds in (2.10), (2.11) apply in this case since  $b > \frac{1}{2}$ . We further note that we can re-estimate  $u$  in  $X_{k,\frac{1}{2}}^S$  in (2.6) to obtain

$$\|u\|_{X_{k,\frac{1}{2}}^S} \lesssim \|u_0\|_{H^k} + (\|n_0\|_{H^s} + \|n_1\|_{H^{s-1}} + \|u_0\|_{H^k}^2) \|u_0\|_{H^k}. \quad (2.12)$$

### 3. WAVE NORM-INFLATION FOR $s > 2k - \frac{1}{2}$

Here we prove Theorem 1.2. In Steps 1-3, the result will be established for  $0 < k < \frac{7}{4}$  and  $s > 2k - \frac{1}{2}$  but with  $s$  near  $2k - \frac{1}{2}$ . In Steps 4-5, the general case of the theorem is reduced to the case considered in Steps 1-3.

*Proof.* Let  $0 < k < 1$ . Let

$$\begin{aligned} \hat{\phi}_{N,A}(\xi) &= N^{\frac{1}{2}-k} \chi_{[-N-\frac{1}{N}, -N]}(\xi) \\ \hat{\phi}_{N,B}(\xi) &= N^{\frac{1}{2}-k} \chi_{[N+1, N+1+\frac{1}{N}]}(\xi) \end{aligned}$$

Let  $\phi_N = \phi_{N,A} + \phi_{N,B}$ . Then  $\|\phi_N\|_{H^k} \sim 1$ . A solution to the integral equation

$$u_N(t) = \psi_1(t)U(t)\phi_N - i\psi_T(t)U *_R \{ [W_+ *_R (\partial_x |u_N|^2) - W_- *_R (\partial_x |u_N|^2)] \cdot u_N \}(t) \quad (3.1)$$

provides a solution to (1.1) with initial data  $(\phi_N, 0, 0)$  when  $n_N$  is defined in terms of  $u_N$  as

$$n_N = W_+ *_R (\partial_x |u_N|^2) - W_- *_R (\partial_x |u_N|^2) \quad (3.2)$$

By working with the estimates in Lemmas 2.3, 2.4 (taking  $s = k - \sigma - 1/2$  in the discussion of §2), we obtain a solution  $u_N$  to (3.1) in  $X_{k-\sigma, \frac{3}{4}-2\epsilon}^S$  for  $0 \leq \sigma \leq k$ , on  $[0, T]$ , where  $T = T(\|\phi_N\|_{H^{k-\sigma}})$  (thus independent of  $N$ ) satisfying

$$\|u_N\|_{C([0,T]; H_x^{k-\sigma})} \leq \|u_N\|_{X_{k-\sigma, \frac{3}{4}-2\epsilon}^S} \leq \|\phi_N\|_{H^{k-\sigma}} \sim N^{-\sigma} \quad (3.3)$$

**Step 1.** We show that

$$\|[(W_+ - W_-) *_R \partial_x |U\phi_N|^2](t)\|_{H^s} \sim tN^{s-(2k-\frac{1}{2})} \quad \text{for } N \gtrsim t^{-1} \quad (3.4)$$

That says that (1.2) holds provided  $u_N(t)$  is replaced by the linear flow  $U(t)\phi_N$  in (3.2).

To show this, note that in the pairing  $U(t)\phi_N \overline{U(t)\phi_N}$ , there are 4 combinations  $U(t)\phi_{N,j} \overline{U(t)\phi_{N,k}}$ , where  $j, k \in \{A, B\}$ . We claim that

$$[W_+ *_R \partial_x (U\phi_{N,A} \overline{U\phi_{N,B}})](\xi, t) \sim i\xi t N^{1-2k} e^{-it\xi} h_1(\xi) \quad (3.5)$$

where  $h_1(\xi)$  is the ‘‘triangular step function’’ with peak at  $\xi = -2N - 1 - \frac{1}{N}$ , of width  $\frac{2}{N}$ , and of height  $\frac{1}{N}$ , i.e.

$$h_1(\xi) = \begin{cases} \xi - (-2N - 1 - \frac{2}{N}) & \text{if } \xi \in [-2N - 1 - \frac{2}{N}, -2N - 1 - \frac{1}{N}] \\ (-2N - 1) - \xi & \text{if } \xi \in [-2N - 1 - \frac{1}{N}, -2N - 1] \end{cases}$$

Here, the symbol  $\sim$  means that the difference between the two quantities has  $H^s$  norm of lower order in  $N$ . It then follows by taking complex conjugates in (3.5) that

$$[W_+ *_R \partial_x(U\phi_{N,B}\overline{U\phi_{N,A}})](t, \xi) \sim i\xi t e^{-it\xi} N^{1-2k} h_2(\xi) \quad (3.6)$$

where  $h_2(\xi)$  is the ‘‘triangular step function’’ centered at  $2N + 1 + \frac{1}{N}$ , of width  $\frac{2}{N}$ , and of height  $\frac{1}{N}$ , i.e.

$$h_2(\xi) = \begin{cases} \xi - (2N + 1) & \text{if } \xi \in [2N + 1, 2N + 1 + \frac{1}{N}] \\ (2N + 1 + \frac{2}{N}) - \xi & \text{if } \xi \in [2N + 1 + \frac{1}{N}, 2N + 1 + \frac{2}{N}] \end{cases}$$

Hence

$$\|[W_+ *_R \partial_x(U\phi_{N,A}\overline{U\phi_{N,B}} + U\phi_{N,B}\overline{U\phi_{N,A}})](t)\|_{H_x^s} \sim tN^{s-(2k-\frac{1}{2})} \quad (3.7)$$

We further claim that the  $AA$  and  $BB$  interactions for the  $W_+$  term are of lower order in  $N$ , i.e. specifically,

$$\|[W_+ *_R \partial_x(U\phi_{N,j}\overline{U\phi_{N,k}})](t)\|_{H^s} \leq N^{s-(2k-\frac{1}{2})-1} \quad \text{for } j = k = A \text{ and } j = k = B \quad (3.8)$$

Finally, we claim that all of the interactions  $AA$ ,  $AB$ ,  $BA$ , and  $BB$  for the  $W_-$  term are of lower order in  $N$ , i.e.

$$\|[W_- *_R \partial_x(U\phi_{N,j}U\phi_{N,k})](t)\|_{H^s} \leq N^{s-(2k-\frac{1}{2})-1} \quad \text{for } j, k \in \{A, B\} \quad (3.9)$$

Combining (3.7), (3.8) (3.9) establishes (3.4). We begin by proving (3.5). Note that

$$\begin{aligned} U(t)\phi_{N,A}(x) &= N^{\frac{1}{2}-k} \int_{\xi_1 \in [-N-\frac{1}{N}, -N]} e^{ix\xi_1} e^{-it\xi_1^2} d\xi_1 \\ \overline{U(t)\phi_{N,B}(x)} &= N^{\frac{1}{2}-k} \int_{\xi_2 \in [-N-1-\frac{1}{N}, -N-1]} e^{ix\xi_2} e^{it\xi_2^2} d\xi_2 \end{aligned}$$

after the change of variable  $\xi_2 \mapsto -\xi_2$  in the second equation. For the remainder of the computation,  $\xi_1$  is restricted to  $[-N - \frac{1}{N}, -N]$  and  $\xi_2$  is restricted to  $[-N - 1 - \frac{1}{N}, -N - 1]$ . By (2.3),

$$\begin{aligned} &W_+ *_R \partial_x(U\phi_{N,A}\overline{U\phi_{N,B}})(t) \\ &= N^{1-2k} \int_{s=0}^t \int_{\xi_1} \int_{\xi_2} i(\xi_1 + \xi_2) e^{i(x-s)(\xi_1+\xi_2)} e^{-i(t-s)(\xi_1^2-\xi_2^2)} d\xi_1 d\xi_2 ds \\ &= N^{1-2k} \int_{\xi_1} \int_{\xi_2} i(\xi_1 + \xi_2) e^{ix(\xi_1+\xi_2)} e^{it(\xi_1^2-\xi_2^2)} g(t, \xi_1, \xi_2) d\xi_1 d\xi_2 \end{aligned}$$

where

$$g(t, \xi_1, \xi_2) = \int_{s=0}^t e^{-is(\xi_1+\xi_2)} e^{is(\xi_1^2-\xi_2^2)} ds = \frac{e^{it(\xi_1+\xi_2)(\xi_1-\xi_2-1)} - 1}{i(\xi_1 + \xi_2)(\xi_1 - \xi_2 - 1)}$$

Since  $\xi_1 + \xi_2$  is confined to a  $\frac{1}{N}$ -sized interval around  $-2N - 1$  and  $\xi_1 - \xi_2 - 1$  is confined to a  $\frac{1}{N}$ -sized interval around 0, we have that  $(\xi_1 + \xi_2)(\xi_1 - \xi_2 - 1)$  is confined to a unit-sized interval around 0. By the power series expansion for  $e^z$ , we

have  $g(t, \xi_1, \xi_2) \sim t$ .

$$\begin{aligned} & [W_+ *_R \partial_x(U\phi_{N,A}\overline{U\phi_{N,B}})(t)](\xi, t) \\ &= N^{1-2k} \int_{\xi_1} \int_{\xi_2} i(\xi_1 + \xi_2)\delta(\xi_1 + \xi_2 - \xi)e^{-it(\xi_1^2 - \xi_2^2)}g(t, \xi_1, \xi_2) d\xi_1 d\xi_2 \end{aligned}$$

Using that  $e^{-it(\xi_1^2 - \xi_2^2)} = e^{-it(\xi_1 - \xi_2 - 1)(\xi_1 + \xi_2)}e^{-it(\xi_1 + \xi_2)} \sim e^{-it(\xi_1 + \xi_2)}$  and that  $g(t, \xi_1, \xi_2) \sim t$ , we obtain (3.5). (3.8) and (3.9) are proved by a similar computation; we only present the proof of (3.8) in the case  $j = k = A$ . For  $t \in [0, T]$ ,

$$W_+ *_R \partial_x(U\phi_{N,A}\overline{U\phi_{N,A}})(t) = \int_{\tau, \xi} \xi e^{ix\xi} \frac{e^{it\tau} - e^{-it\xi}}{\tau + \xi} g(\tau, \xi) d\tau d\xi \tag{3.10}$$

where

$$\begin{aligned} g(\tau, \xi) &= \iint_{\substack{\xi = \xi_1 + \xi_2 \\ \tau = \tau_1 + \tau_2}} [\psi_1 U\phi_{N,A}](\xi_1, \tau_1) [\overline{\psi_1 U\phi_{N,A}}](\xi_2, \tau_2) \\ &= \iint_{\substack{\xi = \xi_1 + \xi_2 \\ \tau = \tau_1 + \tau_2}} \hat{\psi}_1(\tau_1 + \xi_1^2) \hat{\phi}_{N,A}(\xi_1) \hat{\psi}_1(\tau_2 - \xi_2^2) \overline{\hat{\phi}_{N,A}(-\xi_2)} \end{aligned}$$

In this integral,  $\xi_1$  and  $\xi_2$  are each confined to a  $\frac{1}{N}$  sized interval around  $-N$ , forcing  $\xi$  to lie in a  $\frac{1}{N}$  sized interval around  $-2N$ . The  $\hat{\psi}_1(\tau_1 + \xi_1^2)$  and  $\hat{\psi}_1(\tau_2 - \xi_2^2)$  factors then (essentially) restrict  $\tau_1$  to a unit sized interval around  $-N^2$  and restrict  $\tau_2$  to a unit sized interval around  $N^2$ , so that  $\tau$  is forced to lie within a unit sized interval around 0. Consequently,

$$g(\xi, \tau) \begin{cases} \leq \frac{1}{N} & \text{if } (\xi, \tau) \in [-2N - \frac{2}{N}, -2N] \times [-1, 1] \\ = 0 & \text{otherwise} \end{cases}$$

On the support of  $g(\xi, \tau)$ , the factor  $|\tau + \xi| \sim N$ . From (3.10),

$$\begin{aligned} & \| [W_+ *_R \partial_x(U\phi_{N,A}\overline{U\phi_{N,A}})](t) \|_{H^s} \\ & \leq N^{1-2k} \left( \int_{\xi} |\xi|^2 \langle \xi \rangle^{2s} \left[ \int_{\tau} \frac{|g(\tau, \xi)|}{|\tau + \xi|} d\tau \right]^2 d\xi \right)^{1/2} \\ & \leq N^{s - (2k - \frac{1}{2}) - 1} \end{aligned}$$

**Step 2.** Also, on this time interval  $[0, T]$  independent of  $N$ , we claim that

$$\|u_N - \psi_1(t)U(t)\phi_N\|_{X_{k+\sigma, b_1}^S} \leq \|\phi_N\|_{H^{k'}}^2 \|\phi_N\|_{H^{k+\sigma}} \sim N^{2(k'-k)+\sigma} \tag{3.11}$$

where

$$b_1 = \begin{cases} \frac{3}{4} - \frac{k+\sigma}{2} & \text{if } 0 < k + \sigma < \frac{1}{2} \\ \frac{1}{2} & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases} \quad k' = \begin{cases} 0 & \text{if } 0 < k + \sigma < \frac{1}{2} \\ \frac{k+\sigma}{2} - \frac{1}{4} & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases} \tag{3.12}$$

Note that  $2(k' - k) + \sigma$  will be  $< 0$  provided  $\sigma > 0$  is not chosen too large. This says that  $u_N(t)$  is well-approximated by the linear flow  $\psi_1(t)U(t)\phi_N$  in the *stronger* norm  $X_{k+\sigma}^S$ .

We now prove (3.11). From (3.1),

$$\begin{aligned} & \|u_N - \psi_1 U\phi_N\|_{X_{k+\sigma, b_1}^S} \\ & \leq \begin{cases} \| (W_{\pm} *_R \partial_x |u_N|^2) \cdot u_N \|_{X_{k+\sigma, -c_1}^S} & \text{if } 0 \leq k + \sigma \leq \frac{1}{2} \\ \| (W_{\pm} *_R \partial_x |u_N|^2) \cdot u_N \|_{X_{k+\sigma, -c_1}^S \cap Y_{k+\sigma}^S} & \text{if } \frac{1}{2} \leq k + \sigma \leq \frac{5}{2} \end{cases} \end{aligned}$$

for  $b_1$  as defined above and

$$c_1 = \begin{cases} \frac{1}{4} + \frac{k+\sigma}{2} & \text{if } 0 < k + \sigma < \frac{1}{2} \\ \frac{1}{2} & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases}$$

Following with Lemma 2.3,

$$\|u_N - \psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^S} \leq \|W_{\pm} *_R \partial_x |u_N|^2\|_{X_{s', b}^{W_{\pm}}} \|u_N\|_{X_{k+\sigma, b_1}^S}$$

where

$$s' = \begin{cases} -\frac{1}{2} & \text{if } 0 < k + \sigma < \frac{1}{2} \\ k + \sigma - 1 & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases} \quad b = \begin{cases} \frac{1}{2} - \frac{k+\sigma}{2} + \epsilon & \text{if } 0 < k + \sigma < \frac{1}{2} \\ \frac{1}{4} + \epsilon & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases}$$

By Lemma 2.4,

$$\|W_{\pm} *_R \partial_x |u_N|^2\|_{X_{s', b}^{W_{\pm}}} \leq \|\partial_x |u_N|^2\|_{X_{s', -c}^{W_{\pm}}} \leq \|u_N\|_{X_{k', b'_1}^S}^2$$

where

$$c = 1 - b = \begin{cases} \frac{1}{2} + \frac{k+\sigma}{2} - \epsilon & \text{if } 0 < k + \sigma < \frac{1}{2} \\ \frac{3}{4} - \epsilon & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases}$$

$$b'_1 = \begin{cases} \frac{1}{4} + \epsilon & \text{if } 0 < k + \sigma \leq \frac{1}{2} \\ \frac{k+\sigma}{4} + \frac{1}{8} & \text{if } \frac{1}{2} < k + \sigma < \frac{5}{2} \end{cases}$$

and  $k', b_1$  are defined above. Note that  $b'_1 < \frac{3}{4} - 2\epsilon$ . Combining,

$$\begin{aligned} & \|u_N - \psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^S} \\ & \leq \|u_N\|_{X_{k', b'_1}^S}^2 \|u_N\|_{X_{k+\sigma, b_1}^S} \\ & \leq \|u_N\|_{X_{k', b'_1}^S}^2 \|u_N - \psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^S} + \|u_N\|_{X_{k', b'_1}^S}^2 \|\psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^S}, \end{aligned}$$

which by (3.3) is less than or equal to

$$\|\phi_N\|_{H^{k'}}^2 \|u_N - \psi_1(t)U(t)\phi_N\|_{X_{k, b_1}^S} + \|\phi_N\|_{H^{k'}}^2 \|\phi_N\|_{H^{k+\sigma}}$$

Since  $\|\phi_N\|_{H^{k'}} \sim N^{-(k-k')}$ , provided  $N$  is taken large enough and  $k' < k$ , (3.11) will follow.

**Step 3.** Here, we establish

$$\|r_N(t)\|_{H^s} \geq tN^{s-(2k-\frac{1}{2})} \quad \text{for } N \geq t^{-1}$$

if  $0 < k \leq \frac{1}{4}$  and  $2k - \frac{1}{2} < s \leq 4k - 1/2$ , or if  $\frac{1}{4} \leq k < 1$  and  $2k - \frac{1}{2} < s \leq \frac{4}{3}k + \frac{1}{6}$ . To show this, we note that by (3.2) and (3.4), it suffices to show that

$$\|W_{\pm} *_R \partial_x (|u_N|^2 - |\psi_1 U \phi_N|^2)(t)\|_{H^s} \leq 1.$$

Writing

$$|u_N|^2 - |\psi_1 U \phi_N|^2 = |u_N - \psi_1 U \phi_N|^2 + 2 \operatorname{Re} [(u_N - \psi_1 U \phi_N) \overline{\psi_1 U \phi_N}]$$

we see that it suffices to show that

$$\begin{aligned} & \| [W_{\pm} *_R \partial_x |u_N - \psi_1 U \phi_N|^2](t) \|_{H^s} \leq 1 \\ & \| [W_{\pm} *_R \partial_x (u_N - \psi_1 U \phi_N) \overline{\psi_1 U \phi_N}](t) \|_{H^s} \leq 1 \\ & \| [W_{\pm} *_R \partial_x (\psi_1 U \phi_N \cdot \overline{u_N - \psi_1 U \phi_N})](t) \|_{H^s} \leq 1 \end{aligned} \tag{3.13}$$

We focus on the middle estimate (3.13); the other two are handled similarly. As we describe in detail below, by requiring  $s$  to lie sufficiently close to (but above)  $2k - \frac{1}{2}$ , we can assign  $\sigma > 0$  such that

$$s \leq \begin{cases} 2(k + \sigma) - \frac{1}{2} & \text{if } 0 < k + \sigma < \frac{1}{2} \\ k + \sigma & \text{if } \frac{1}{2} \leq k + \sigma < \frac{5}{2} \end{cases} \tag{3.14}$$

and

$$k' + (k + \sigma) \leq 2k \tag{3.15}$$

where  $k'$  is given in (3.12). Then proceed to estimate the left-hand side of (3.13) by Lemma 2.2(2) as

$$\|u_N - \psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^s} \|\psi_1 U \phi_N\|_{X_{k+\sigma, b_1}^s}$$

By Step 2 and Lemma 2.1(1), the above expression is less than or equal to

$$\|\phi_N\|_{H^{k'}}^2 \|\phi_N\|_{H^{k+\sigma}}^2 \sim N^{2(k'-k)} N^{2(k+\sigma-k)}.$$

By (3.15), it follows that the exponent is less than or equal to zero.

We now provide the details assigning  $\sigma$  in terms of  $k$  and  $s$ . The condition (3.15) is equivalent to the restriction

$$\sigma \leq \begin{cases} k & \text{if } k \leq \frac{1}{4} \\ \frac{1}{3}k + \frac{1}{6} & \text{if } \frac{1}{4} \leq k \end{cases} \tag{3.16}$$

The following assignments meet the criteria (3.16) and (3.14).

- If  $0 < k \leq \frac{1}{4}$ , restrict to  $s$  such that  $2k - \frac{1}{2} < s \leq 4k - 1/2$ , and set  $\sigma = k$ .
- If  $\frac{1}{4} \leq k < 1$ , then restrict to  $s$  such that  $2k - \frac{1}{2} < s \leq \frac{4}{3}k + \frac{1}{6}$  and set  $\sigma = \frac{1}{3}k + \frac{1}{6}$ .

**Step 4.** Suppose  $0 < k < 1$  and  $s > 2k - \frac{1}{2}$ . Let  $s'$  be such that  $s' \leq s$  and  $s'$  meets the restrictions outlined in Step 3 with  $s$  replaced by  $s'$ . Then by Steps 1–3 (with  $s$  replaced by  $s'$ )

$$\|n_N(t)\|_{H^s} \geq \|n_N(t)\|_{H^{s'}} \geq t N^{s' - (2k - \frac{1}{2})} \quad \text{for } N \geq t^{-1}$$

so we can take  $\alpha = s' - (2k - \frac{1}{2})$  in the statement of the theorem.

**Step 5.** Next, suppose  $k < 0$  and  $s > -1/2$ . By the reasoning of Step 4, it suffices to restrict to  $s < \frac{3}{2}$ . Set  $0 < k'' < \frac{1}{2}s + \frac{1}{4}$ , and note that  $s > 2k'' - 1/2$ . Clearly  $\|u_N(t)\|_{H^k} \leq \|u_N(t)\|_{H^{k''}}$ , so we can just appeal to the conclusion of Steps 1–4 applied with  $k$  replaced by  $k''$ .  $\square$

#### 4. A PRELIMINARY ANALYSIS FOR $s \leq -\frac{3}{2}$

Let  $f(x) = \sqrt{2} \operatorname{sech}(x)$ , which is the unique positive ground state solution to

$$-f + \partial_x^2 f + |f|^2 f = 0 \tag{4.1}$$

Let  $f_\lambda(x) = \lambda f(\lambda x)$  and set

$$\begin{aligned} u_{\lambda, N}(x, t) &= e^{it(\lambda^2 - N^2)} e^{iN x} \sqrt{1 - 4N^2} f_\lambda(x - 2Nt) \\ n_{\lambda, N}(x, t) &= -|f_\lambda(x - 2Nt)|^2 \end{aligned}$$

From (4.1), it follows that  $(u_{\lambda, N}, n_{\lambda, N})$  solves (1.1) for all  $\lambda \in \mathbb{R}$  and  $-\frac{1}{2} < N < \frac{1}{2}$ . This is the exact soliton class appearing in [8] and [11].

Our next goal is to prove Theorem 1.3 demonstrating phase decoherence ill-posedness for  $k = 0, s < -\frac{3}{2}$ . We first, however, settle for a partial result (Proposition 4.1) using a pair from the above exact explicit soliton class. We include this result since it is clear and straightforward and exhibits the idea behind the proof of the full result (Theorem 1.3), which is considerably more technical and appears in the next section.

Define the norm for  $H^s(|\xi| \geq M)$  as

$$\|\phi\|_{H^s(|\xi| \geq M)} = \left( \int_{|\xi| \geq M} |\xi|^{2s} |\hat{\phi}(\xi)|^2 d\xi \right)^{1/2}$$

The limitation of the following partial result is the use of  $H^s(|\xi| \geq M)$  and  $H^{s-1}(|\xi| \geq M)$  norms as opposed to the full  $H^s$  and  $H^{s-1}$  norms.

**Proposition 4.1.** *Suppose  $s \leq -\frac{3}{2}$ . Fix any  $T > 0, \delta > 0$ . Then there exist  $M(\delta)$  sufficiently large and  $N(\delta) < \frac{1}{2}$  sufficiently close to  $\frac{1}{2}$  so that if*

$$\lambda_1 = M, \quad \lambda_2 = \sqrt{M^2 + \frac{\pi}{2T}}$$

then the solutions are of unit size on  $[0, T]$ ,

$$\begin{aligned} \|u_{\lambda_j, N}(\cdot, t)\|_{L^2_x} &\sim 1 \\ \|n_{\lambda_j, N}(\cdot, t)\|_{H^s(|\xi| \geq M)} &\sim 1, \quad \|\partial_t n_{\lambda_j, N}(\cdot, t)\|_{H^{s-1}(|\xi| \geq M)} \sim 1 \end{aligned} \tag{4.2}$$

and are initially close

$$\|u_{\lambda_2, N}(\cdot, 0) - u_{\lambda_1, N}(\cdot, 0)\|_{L^2} \leq \delta \tag{4.3}$$

$$\begin{aligned} \|n_{\lambda_2, N}(\cdot, 0) - n_{\lambda_1, N}(\cdot, 0)\|_{H^s(|\xi| \geq M)} &\leq \delta \\ \|\partial_t n_{\lambda_2, N}(\cdot, 0) - \partial_t n_{\lambda_1, N}(\cdot, 0)\|_{H^{s-1}(|\xi| \geq M)} &\leq \delta \end{aligned} \tag{4.4}$$

but become fully separated in the  $u$ -variable by time  $T$ ,

$$\|u_{\lambda_2, N}(\cdot, T) - u_{\lambda_1, N}(\cdot, T)\|_{L^2} \sim 1 \tag{4.5}$$

*Proof.* We will select  $M = M(\delta)$  sufficiently large later. Take  $0 \leq N < \frac{1}{2}$  sufficiently close to  $\frac{1}{2}$  so that  $(1 - 2N)^{1/2} M^{1/2} = 1$ . Then since  $N \sim \frac{1}{2}$  we have  $\sqrt{1 - 4N^2} \sim (1 - 2N)^{1/2}$  and noting that  $\lambda_1 = M$  and  $(1 - 2N)^{1/2} M^{1/2} = 1$  gives

$$\begin{aligned} \|u_{\lambda_2, N}(\cdot, 0) - u_{\lambda_1, N}(\cdot, 0)\|_{L^2} &= (1 - 2N)^{1/2} \left\| \hat{f}\left(\frac{\xi}{\lambda_2}\right) - \hat{f}\left(\frac{\xi}{\lambda_1}\right) \right\|_{L^2_\xi} \\ &= \left\| \hat{f}\left(\frac{\lambda_1 \xi}{\lambda_2}\right) - \hat{f}(\xi) \right\|_{L^2_\xi} \end{aligned}$$

Take  $M$  sufficiently large so that  $\lambda_1/\lambda_2$  is sufficiently close to 1 in order to make the above expression  $\leq \delta$ . Thus (4.3) is established. Next, we establish (4.4). By the change of variable  $\xi \mapsto \lambda_1 \xi$

$$\begin{aligned} &\|n_{\lambda_2, N}(\cdot, 0) - n_{\lambda_1, N}(\cdot, 0)\|_{H^s(|\xi| \geq M)}^2 \\ &= \lambda_1^{3+2s} \int_{|\xi| \geq 1} \left| \frac{\lambda_2}{\lambda_1} (f^2) \left( \frac{\xi \lambda_1}{\lambda_2} \right) - (f^2)(\xi) \right|^2 |\xi|^{2s} d\xi \end{aligned}$$

Since  $s \leq -\frac{3}{2}$  we have  $\lambda_1^{3+2s} \leq 1$  and the above difference is made  $\leq \delta$  by again taking  $M$  sufficiently large. Also

$$\begin{aligned} & \|\partial_t n_{\lambda_2, N}(\cdot, 0) - \partial_t n_{\lambda_1, N}(\cdot, 0)\|_{H^{s-1}(|\xi| \geq M)}^2 \\ &= N^2 \lambda_1^{3+2s} \int_{|\xi| \geq 1} \left| \frac{\lambda_2^2}{\lambda_1^2} (f^2)'(\frac{\xi \lambda_1}{\lambda_2}) - (f^2)'(\xi) \right|^2 |\xi|^{2(s-1)} d\xi \end{aligned}$$

(Here, the notation  $'$  indicates the derivative). Since  $s \leq -\frac{3}{2}$  we have  $\lambda_1^{3+2s} \leq 1$  and the above difference is made  $\leq \delta$  by again taking  $M$  sufficiently large. The statements (4.2) are proved by similar change of variable calculations. The need for the restrictions to  $|\xi| \geq M$  in (4.4) is clear from these calculations. In fact, one can show that for  $s < -\frac{1}{2}$ , we have  $\|n_{\lambda, N}(\cdot, 0)\|_{H^s} \sim \lambda$  as  $\lambda \rightarrow +\infty$  due to the  $|\xi| \leq \lambda$  frequency contribution.

Now we establish (4.5). The key observation here is that while  $\lambda_2 - \lambda_1$  is very small (as  $M \rightarrow +\infty$ ),  $\lambda_2^2 - \lambda_1^2$  is of fixed size  $\pi/(2T)$  and thus  $e^{iT(\lambda_2^2 - \lambda_1^2)} = i$  is purely imaginary. Now

$$\|u_2(\cdot, T) - u_1(\cdot, T)\|_{L^2}^2 = \|u_2(\cdot, T)\|_{L^2}^2 + \|u_1(\cdot, T)\|_{L^2}^2 - 2 \operatorname{Re} \int_x u_2(x, T) \overline{u_1(x, T)} dx$$

but the last term on the right-hand side is

$$-2 \operatorname{Re} e^{iT(\lambda_2^2 - \lambda_1^2)} (1 - 4N^2) \int_x \lambda_2 f(\lambda_2 x) \lambda_1 f(\lambda_1 x) dx = 0$$

which, combined with (4.2) gives (4.5). □

### 5. SCHRÖDINGER PHASE DECOHERENCE FOR $s < -3/2$

Here, we remove the shortcoming of Proposition 4.1 (high frequency truncated norms  $H^s(|\xi| \geq M)$ ,  $H^{s-1}(|\xi| \geq M)$  used instead of  $H^s$ ,  $H^{s-1}$ ) and prove Theorem 1.3. The soliton class employed in the proof of Proposition 4.1 involved assigning

$$n(x, t) = -\lambda^2 |f|^2(\lambda(x - 2tN))$$

and thus  $\hat{n}(\xi, t) = -\lambda(|f|^2)'(\xi/\lambda) e^{-2itN\xi}$ . Replace  $|f|^2$  in the definition of  $n$  by  $g$  defined by  $\hat{g}(\xi) = (|f|^2)'(\xi) \chi_{|\xi| \geq 1}(\xi)$  (i.e. the restriction to frequencies  $\geq 1$ ) and set

$$\tilde{n}(x, t) = -\lambda^2 g(\lambda(x - 2tN))$$

Then

$$\|\tilde{n}(\cdot, t)\|_{H^s} + \|\partial_t \tilde{n}(\cdot, t)\|_{H^{s-1}} \leq 1, \quad \text{as } \lambda \rightarrow +\infty$$

Unfortunately,  $(u, \tilde{n})$  is no longer a solution to (1.1). We shall thus adapt the method of Christ-Colliander-Tao [5] to construct a “near soliton” class that grants more flexibility in the selection of the wave initial data. The method proceeds by solving a “small dispersion approximation” to the equation, and by introducing scaling and phase translation parameters, building the “near soliton” class. The main new obstacle, in comparison to the work of [5] applied to the nonlinear Schrödinger equation, is that (1.1) does not possess scaling nor the Galilean (phase shift) identity. We thus need to carry out the small dispersion approximation for a *modified* Zakharov system with the property that when scaling and phase shift operations are performed, the modified Zakharov system is converted into the true Zakharov system.

Consider fixed initial data  $(n_0, u_0)$  (to be defined later).

**Step 1.** The solution to the small dispersion approximation

$$i\partial_t v = n_0(x)v$$

with  $v(x, 0) = u_0(x)$  is

$$v(x, t) = e^{-itn_0(x)}u_0(x)$$

**Step 2.** For parameters  $\lambda \gg 1$ ,  $0 < \nu \ll 1$ ,  $-\frac{1}{2} < N < \frac{1}{2}$ , consider the initial-value problem for the modified Zakharov system  $u = u^{(\lambda, \nu, N)}$ ,  $n_{\pm} = n_{\pm}^{(\lambda, \nu, N)}$ ,

$$\begin{aligned} i\partial_t u + \nu^2 \partial_x^2 u &= \frac{1}{2} \left[ n_0 \left( x + \frac{\nu(1+2N)}{\lambda} t \right) + n_0 \left( x - \frac{\nu(1-2N)}{\lambda} t \right) \right] u + (n_+ + n_-)u \\ \frac{\lambda}{(1-2N)\nu} \partial_t n_+ + \partial_x n_+ &= -\frac{1}{2}(1+2N)\partial_x |u|^2 \\ \frac{\lambda}{(1+2N)\nu} \partial_t n_- - \partial_x n_- &= \frac{1}{2}(1-2N)\partial_x |u|^2 \\ u(x, 0) &= u_0(x), \quad n_{\pm}(x, 0) = 0 \end{aligned} \tag{5.1}$$

If  $k \geq 1$  and (implicit constants here depend on  $\|u_0\|_{H^k}$  and  $\|n_0\|_{H^k}$ )

$$T \lesssim |\ln \nu|, \quad \lambda \gtrsim \nu^{-5} \tag{5.2}$$

then, on  $[0, T]$ , we have the two estimates

$$\|u\|_{L_T^\infty H_x^k} \lesssim \nu^{-1/2} \tag{5.3}$$

$$\|n_{\pm}\|_{L_T^\infty H_x^{k-1}} \lesssim \frac{1-4N^2}{\lambda} \tag{5.4}$$

To show this, note first that

$$\begin{aligned} n_+ &= -\frac{1}{2}(1+2N) \int_0^{t\mu_+} \partial_x |u|^2 \left( x - s, t - \frac{s}{\mu_+} \right) ds \\ n_- &= \frac{1}{2}(1-2N) \int_0^{t\mu_-} \partial_x |u|^2 \left( x + s, t - \frac{s}{\mu_-} \right) ds \end{aligned}$$

where  $\mu_+ = \nu(1-2N)/\lambda$  and  $\mu_- = \nu(1+2N)/\lambda$ , and thus for  $k \geq 1$ ,

$$\|n_{\pm}\|_{L_T^\infty H_x^{k-1}} \leq \frac{(1-4N^2)\nu T}{\lambda} \|u\|_{L_T^\infty H_x^k}^2 \tag{5.5}$$

By the energy method applied to (5.1), we have

$$\begin{aligned} &\|\partial_x^k u(T)\|_{L_x^2}^2 - \|\partial_x^k u(0)\|_{L_x^2}^2 \\ &= -\operatorname{Re} i \int_0^T \int_x \partial_x^k [n_0(\dots)u] \overline{\partial_x^k u} \, dx \, dt - 2 \operatorname{Re} i \int_0^T \int_x \partial_x^k [n_{\pm}u] \overline{\partial_x^k u} \, dx \, dt \\ &= \text{I} + \text{II} \end{aligned}$$

Term I will be addressed via the Gronwall inequality, while in estimating II we will produce a small coefficient.

$$|\text{I}| \leq \|n_0\|_{H^k} \int_0^T \|u(t)\|_{H_x^k}^2 \, dt$$

Term II is decomposed as

$$\begin{aligned} \text{II} &= -\operatorname{Re} i \sum_{\substack{\alpha+\beta=k \\ \alpha \leq k-1}} c_{\alpha\beta} \int_0^T \int_x \partial_x^\alpha n_\pm \partial_x^\beta u \overline{\partial_x^k u} \, dx \, dt - 2 \operatorname{Re} i \int_0^T \int_x \partial_x^k n_\pm u \overline{\partial_x^k u} \, dx \, dt \\ &= \text{II}_a + \text{II}_b \end{aligned}$$

From (5.5),

$$|\text{II}_a| \leq T \|n_\pm\|_{L_T^\infty H_x^{k-1}} \|u\|_{L_T^\infty H_x^k}^2 \leq \frac{(1-4N^2)\nu T^2}{\lambda} \|u\|_{L_T^\infty H_x^k}^4 \tag{5.6}$$

while for  $\text{II}_b$ , we integrate by parts (here  $\approx$  means up to terms bounded similarly to (5.6))

$$\begin{aligned} \text{II}_b &= 2 \operatorname{Re} i \int_0^T \int_x \partial_x^{k-1} n_\pm \partial_x [u \overline{\partial_x^k u}] \, dx \, dt \\ &\approx \int_0^T \int_x \partial_x^{k-1} n_\pm \partial_x^{k-1} (iu \overline{\partial_x^2 u} - i\bar{u} \partial_x^2 u) \, dx \, dt \\ &= -\frac{1}{\nu^2} \int_0^T \int_x \partial_x^{k-1} n_\pm \partial_x^{k-1} \partial_t |u|^2 \, dx \, dt \\ &= -\frac{1}{\nu^2} \int_x \partial_x^{k-1} n_\pm(T) \partial_x^{k-1} |u|^2(T) \, dx \, dt + \frac{1}{\nu^2} \int_0^T \int_x \partial_t \partial_x^{k-1} n_\pm \partial_x^{k-1} |u|^2 \, dx \, dt \\ &= \text{II}_{b1} + \text{II}_{b2} \end{aligned}$$

From (5.5), we have

$$|\text{II}_{b1}| \leq \frac{(1-4N^2)T}{\lambda\nu} \|u\|_{L_T^\infty H_x^k}^4$$

From (5.1),  $\partial_x^{k-1} \partial_t n_\pm = \mp \mu_\pm \partial_x^k n_\pm \mp \frac{1}{2} \mu_\pm (1 \pm 2N) \partial_x^k |u|^2$ , so

$$\begin{aligned} \text{II}_{b2} &= \pm \frac{\mu_\pm}{\nu^2} \int_0^T \int_x \partial_x^{k-1} n_\pm \partial_x^k |u|^2 \, dx \, dt \\ \implies |\text{II}_{b2}| &\leq \frac{(1-4N^2)T^2}{\lambda^2} \|u\|_{L_T^\infty H_x^k}^4 \end{aligned}$$

All together, (using  $L^2$  conservation as well),

$$\|u(T)\|_{H_x^k}^2 \leq \|u_0\|_{H^k}^2 + \|n_0\|_{H^k} \int_0^T \|u(t)\|_{H_x^k}^2 \, dt + \epsilon \|u\|_{L_T^\infty H_x^k}^4$$

with

$$\epsilon = \frac{(1-4N^2)T}{\lambda} \left( \frac{T}{\lambda} + \frac{1}{\nu} + \nu T \right) \leq \frac{1}{\lambda^{1/2}}$$

where the last inequality follows from the assumptions (5.2). By the Gronwall inequality,

$$\|u\|_{L_T^\infty H_x^k}^2 \leq e^{\|n_0\|_{H^k} T} (\|u_0\|_{H_x^k}^2 + \lambda^{-1/2} \|u\|_{L_T^\infty H_x^k}^4) \tag{5.7}$$

Provided that we have

$$\lambda^{-1/2} \lesssim e^{-\|n_0\|_{H^k} T} \|u_0\|_{H^k}^{-2}, \tag{5.8}$$

we obtain, from (5.7) and a continuity in time argument, the bound

$$\|u\|_{L_T^\infty H_x^k}^2 \leq 2e^{\|n_0\|_{H^k} T} \|u_0\|_{H^k}^2 \tag{5.9}$$

Now, the assumptions (5.2) imply (5.7); and (5.9) implies (5.3) by the first of the assumptions in (5.2).

**Step 3.** With  $u = u^{(\lambda, \nu, N)}$  as defined in Step 2,  $v$  as defined in Step 1, and (5.2) satisfied, we claim that

$$\|u - v\|_{L_T^\infty H_x^k} \lesssim \nu$$

where the implicit constant depends on  $\|u_0\|_{H_x^{k+2}}$  and  $\|n_0\|_{H_x^{k+2}}$ . For this, we appeal to the result of Step 2 at the level of  $k + 2$  derivatives, and then apply the energy method to the difference  $u - v$  in  $H^k$ :

$$\begin{aligned} & \|\partial_x^k(u - v)(T)\|_{L_x^2}^2 \\ &= -2 \operatorname{Re} i \nu^2 \int_0^T \int_x \partial_x^{k+2} u \overline{\partial_x^k(u - v)} dx dt \\ & \quad - 2 \operatorname{Re} i \int_0^T \int_x \partial_x^k \left[ n_0 \left( x \pm \frac{\nu(1 \pm 2N)}{\lambda} t \right) u - n_0(x)v \right] \overline{\partial_x^k(u - v)} dx dt \\ & \quad - 2 \operatorname{Re} i \int_0^T \int_x \partial_x^k [n_\pm u] \overline{\partial_x^k(u - v)} dx dt \\ &= \text{I} + \text{II} + \text{III} \end{aligned}$$

Direct estimates using (5.3), (5.4) give

$$|\text{I}| \leq T^2 \nu^4 \|u\|_{L_T^\infty H_x^{k+2}}^2 + \frac{1}{4} \|\partial_x^k(u - v)\|_{L_T^\infty L_x^2}^2 \leq cT^2 \nu^3 + \frac{1}{4} \|\partial_x^k(u - v)\|_{L_T^\infty L_x^2}^2.$$

By (5.3) and (5.4),

$$\begin{aligned} |\text{III}| &\leq T^2 \|n_\pm\|_{L_T^\infty H_x^k}^2 \|u\|_{L_T^\infty H_x^k}^2 + \frac{1}{4} \|\partial_x^k(u - v)\|_{L_T^\infty L_x^2}^2 \\ &\leq \frac{c(1 - 4N^2)^2 T^2}{\lambda^2 \nu} + \frac{1}{4} \|\partial_x^k(u - v)\|_{L_T^\infty L_x^2}^2 \end{aligned}$$

By rewriting

$$\begin{aligned} & n_0 \left( x \pm \frac{\nu(1 \pm 2N)}{\lambda} t \right) u - n_0(x)v \\ &= \left[ \int_0^{\pm \frac{\nu(1 \pm 2N)t}{\lambda}} \partial_x n_0(x + s) ds \right] u + n_0(x)(u - v) \end{aligned}$$

term II can be estimated as

$$|\text{II}| \leq \frac{\nu(1 \pm 2N)T^2}{\lambda} \|n_0\|_{H^{k+1}} \|u\|_{L_T^\infty H_x^k} \|u - v\|_{L_T^\infty H_x^k} + \|n_0\|_{H_x^k} \int_0^T \|(u - v)(t)\|_{H_x^k}^2 dt$$

Combining, and applying the Gronwall inequality, we have

$$\|u - v\|_{L_T^\infty H_x^k}^2 \lesssim e^{cT} \left[ \frac{\nu(1 \pm 2N)^2 T^4}{\lambda^2} + T^2 \nu^3 + \frac{(1 - 4N^2)^2 T^2}{\lambda^2 \nu} \right]$$

The result follows from the assumptions (5.2).

**Step 4.** For  $-\frac{1}{2} < N < \frac{1}{2}$ , set

$$\begin{aligned} u(x, t) &= \lambda(1 - 4N^2)^{1/2} e^{ixN} e^{-itN^2} u^{(\lambda, \nu, N)}(\lambda\nu(x - 2tN), \lambda^2 t) \\ n_\pm(x, t) &= \lambda^2 n_\pm^{(\lambda, \nu, N)}(\lambda\nu(x - 2tN), \lambda^2 t). \end{aligned}$$

Then  $(u, n)$  solves (1.1), with

$$n(x, t) = n_+(x, t) + n_-(x, t) + \frac{1}{2} \lambda^2 n_0(\lambda\nu(x + t)) + \frac{1}{2} \lambda^2 n_0(\lambda\nu(x - t))$$

and initial data  $u(x, 0) = \lambda u_0(\lambda \nu x)$ ,  $n(x, 0) = \lambda^2 n_0(\lambda \nu x)$ ,  $\partial_t n(x, 0) = 0$ .

Consider  $0 < \nu \ll 1$  and  $\lambda \gg 1$ . Since  $\|u^{(\lambda, \nu, N)}(x, t)\|_{L_x^2} = \|u_0\|_{L_x^2}$  for all  $t$ , we have by change of variable

$$\|u(x, t)\|_{L_x^2} = \frac{\lambda^{1/2}(1 - 4N^2)^{1/2}}{\nu^{1/2}} \|u_0\|_{L_x^2}$$

Also, if  $\hat{n}_0(\xi) = 0$  for  $|\xi| \leq 1$  and  $\lambda \nu \geq 1$ , then another change of variable gives

$$\|n(x, 0)\|_{H_x^s} \leq \lambda^{\frac{3}{2}+s} \nu^{s-\frac{1}{2}} \|n_0\|_{H_x^s}$$

If  $s < -3/2$  and  $\lambda$  and  $\nu$  satisfy

$$\lambda \geq \nu^{-\alpha}, \quad \text{with } \alpha = \max\left(\frac{\frac{1}{2} - s}{-s - \frac{3}{2}}, 5\right) \tag{5.10}$$

then  $\|n(x, 0)\|_{H_x^s} \leq \|n_0\|_{H_x^s}$ .

**Step 5.** Fix  $M \gg 1$  and  $0 \leq \nu \ll 1$ , to be chosen momentarily. In terms of  $M$  and  $\nu$ , define the following quantities: Let  $T = |\ln \nu|/M^2$ , and set

$$\lambda_1 = M, \quad \lambda_2 = \sqrt{\frac{\pi}{2T} + M^2}$$

We note that

$$e^{iT(\lambda_2^2 - \lambda_1^2)} = i \tag{5.11}$$

is purely imaginary. Note further that

$$\frac{\lambda_2}{\lambda_1} = \sqrt{\frac{\pi}{2|\ln \nu|} + 1} \rightarrow 1, \quad \text{as } \nu \rightarrow 0, \quad \text{independently of } M \tag{5.12}$$

Take  $N$  sufficiently close to  $1/2$  so that

$$\frac{(1 - 2N)M}{\nu} = 1 \tag{5.13}$$

Take  $u_0(x) \in \mathcal{S}(\mathbb{R})$  such that  $u_0(x) = 1$  for  $-1 \leq x \leq 1$ , and  $n_0(x)$  to be the smooth function given on the Fourier side as

$$\hat{n}_0(\xi) = \begin{cases} 0 & \text{if } |\xi| \leq 2 \\ \pi/2 & \text{if } 2 \leq |\xi| \leq 4 \\ 0 & \text{if } |\xi| \geq 4 \end{cases}$$

so that in fact

$$n_0(x) = \frac{\cos 3x \sin x}{x}$$

Now consider the solutions  $u^{(\lambda_1, \nu, N)}$  and  $u^{(\lambda_2, \nu, N)}$  of the modified Zakharov system given at the beginning of Step 2, both in terms of the data  $u_0(x)$  and  $n_0(x)$ . Define, as in Step 4, the (1.1) solution  $(u_1, n_1)$  in terms of  $u^{(\lambda_1, \nu, N)}$  and  $(u_2, n_2)$  in terms of  $u^{(\lambda_2, \nu, N)}$ . By the comments at the end of Step 4,

$$\begin{aligned} \|u_j(x, t)\|_{L_x^2} &= 1, \\ \|n_j(x, 0)\|_{H_x^s} + \|\partial_t n_j(x, 0)\|_{H_x^{s-1}} &\leq 1 \end{aligned}$$

where, in order to meet condition (5.10), we need  $M = M(\nu) \geq \nu^{-\alpha}$ . By a change of variable and (5.13),

$$\|u_2(x, t) - u_1(x, t)\|_{L_x^2} = \left\| \frac{\lambda_2}{\lambda_1} u^{(\lambda_2, \nu, N)}\left(\frac{\lambda_2 x}{\lambda_1}, \lambda_2^2 t\right) - u^{(\lambda_1, \nu, N)}(x, \lambda_1^2 t) \right\|_{L_x^2}$$

By (5.12) and the fact that  $\|u^{(\lambda,\nu,N)}(x,t)\|_{L_x^2} = \|u_0\|_{L_x^2}$  for all  $t$  and uniformly in all the parameters, we can take  $\nu = \nu(\delta) > 0$  sufficiently small so that

$$\|u_2(x,t) - u_1(x,t)\|_{L_x^2} = \|u^{(\lambda_2,\nu,N)}(x,\lambda_2^2 t) - u^{(\lambda_1,\nu,N)}(x,\lambda_1^2 t)\|_{L_x^2} + \mathcal{O}(\delta)$$

By the results of Step 1 and 3, again taking  $\nu = \nu(\delta)$  sufficiently small, if  $0 \leq t \leq |\ln \nu|/M^2$ , then

$$\|u_2(x,t) - u_1(x,t)\|_{L_x^2} = \|(e^{i(\lambda_2^2 - \lambda_1^2)t n_0(x)} - 1)u_0(x)\|_{L_x^2} + \mathcal{O}(\delta) \tag{5.14}$$

Here, the first condition of (5.2) is met, since the  $T$  appearing there is our  $\lambda_1^2 t \sim \lambda_2^2 t \lesssim |\ln \nu|$ . We see trivially from (5.14) that

$$\|u_2(x,0) - u_1(x,0)\|_{L_x^2} \lesssim \delta$$

But by (5.11) and (5.14) and the choice of  $u_0(x)$  and  $n_0(x)$ ,

$$\|u_2(x,T) - u_1(x,T)\|_{L_{[0,T]}^\infty L_x^2} = \mathcal{O}(1)$$

We further note that

$$T = \frac{|\ln \nu|}{M^2} \leq |\ln \nu| \nu^4 \rightarrow 0 \quad \text{as } \nu \rightarrow 0$$

and therefore we can accommodate an arbitrarily small preselected time,  $T$  as in the statement of the theorem.

### 6. THE SCHRÖDINGER FLOW MAP IS NOT $C^2$ FOR $s < -1/2$

In this section, we give the proof of Theorem 1.4. For fixed  $H^\infty$  data  $(u_0, n_0, n_1)$ , to be specified later, and a parameter  $\gamma \in \mathbb{R}$ . We consider the initial data

$$(u|_{t=0}, n|_{t=0}, \partial_t n|_{t=0}) = (\gamma u_0, \gamma n_0, \gamma n_1)$$

and the corresponding 1DZS solutions  $(u, n) = (u_\gamma, n_\gamma)$ . Clearly

$$u|_{\gamma=0} = 0, \quad \partial_x u|_{\gamma=0} = 0, \quad \partial_x^2 u|_{\gamma=0} = 0, \quad n|_{\gamma=0} = 0, \quad \partial_x n|_{\gamma=0} = 0 \tag{6.1}$$

The solution, written in integral equation form, is:

$$\begin{aligned} u(t) &= U(t)(\gamma u_0) - i \int_0^t U(t-t')[(un)(t')] dt' \\ n(t) &= W(t)(\gamma n_0, \gamma n_1) \pm \frac{1}{2} \int_0^t \partial_x |u|^2(x \pm s, t-s) ds \end{aligned}$$

from which it follows that

$$\begin{aligned} \partial_\gamma u(t) &= U(t)u_0 - i \int_0^t U(t-t')[(\partial_\gamma u n + u \partial_\gamma n)(t')] dt' \\ \partial_\gamma n(t) &= W(t)(n_0, n_1) \pm \frac{1}{2} \int_0^t \partial_x (\partial_\gamma u \bar{u} + u \overline{\partial_\gamma u})(x \pm s, t-s) ds \end{aligned} \tag{6.2}$$

By (6.1),

$$\partial_\gamma u|_{\gamma=0} = Uu_0, \quad \partial_\gamma n|_{\gamma=0} = W(n_0, n_1) \tag{6.3}$$

By applying  $\partial_x$  to (6.2) and again appealing to (6.1), we get

$$\partial_x \partial_\gamma u|_{\gamma=0} = \partial_x Uu_0, \quad \partial_\gamma \partial_x n|_{\gamma=0} = \partial_x W(n_0, n_1) \tag{6.4}$$

By applying  $\partial_\gamma$  to (6.2), we obtain

$$\begin{aligned} \partial_\gamma^2 u(t) &= -i \int_0^t U(t-t')[(\partial_\gamma^2 u n + 2\partial_\gamma u \partial_\gamma n + u \partial_\gamma^2 n)(t')] dt' \\ \partial_\gamma^2 n(t) &= \pm \int_0^t \partial_x(\partial_\gamma^2 u \bar{u} + 2|\partial_\gamma u|^2 + u \overline{\partial_\gamma^2 u})(x \pm s, t-s) ds \end{aligned}$$

from which we find, together with (6.3)(6.4), that

$$\begin{aligned} \partial_\gamma^2 u|_{\gamma=0}(t) &= -2i \int_0^t U(t-t')[U u_0(t') W(n_0, n_1)(t')] dt' \\ \partial_\gamma^2 n|_{\gamma=0}(t) &= \pm \int_0^t \partial_x |U(t') u_0|^2(x \mp s, t-s) ds \end{aligned}$$

Let  $X = H^k \times H^s \times H^{s-1}$ ,  $Y = H^k \times H^s$ . Fix  $t > 0$ , and let  $F : X \rightarrow Y$  be the solution map  $F(u_0, n_0, n_1) = (u(t), n(t))$ . Let  $G : \mathbb{R} \rightarrow X$  be given by  $G(\gamma) = (\gamma u_0, \gamma n_0, \gamma n_1)$ . Let  $H(\gamma) = F \circ G(\gamma)$  so that  $H : \mathbb{R} \rightarrow Y$ . Then (here  $\mathcal{L}(A; B)$  denotes a linear map  $A \rightarrow B$ )

$$\underbrace{DH(\gamma)}_{\mathcal{L}(\mathbb{R}; Y)} = \underbrace{DF(G(\gamma))}_{\mathcal{L}(X; Y)} \circ \underbrace{DG(\gamma)}_{\mathcal{L}(\mathbb{R}; X)}$$

Also

$$D^2 H(\gamma) = \underbrace{D^2 F(G(\gamma))}_{\mathcal{B}(X \times X; Y)} \circ \underbrace{(DG(\gamma), DG(\gamma))}_{\mathcal{L}(\mathbb{R}; X)} + \underbrace{DF(G(\gamma))}_{\mathcal{L}(X; Y)} \circ \underbrace{D^2 G(\gamma)}_{\mathcal{B}(\mathbb{R} \times \mathbb{R}; X)}$$

and since  $D^2 G(\gamma) = 0$ ,

$$D^2 H(\gamma)(\alpha_1, \alpha_2) = D^2 F(G(\gamma))((\alpha_1 u_0, \alpha_1 n_0, \alpha_1 n_1), (\alpha_2 u_0, \alpha_2 n_0, \alpha_2 n_1))$$

Hence

$$D^2 F(0)((u_0, n_0, n_1), (u_0, n_0, n_1)) = D^2 H(0)(1, 1) = (\partial_\gamma^2 u|_{\gamma=0}(t), \partial_\gamma^2 n|_{\gamma=0}(t))$$

We now note how to prescribe an appropriate sequence  $(u_{N,0}, n_{N,0}, n_{N,1})$  (indexed by  $N$ ) to show that  $D^2 F(0) \in \mathcal{B}(X \times X; Y)$  is not a bounded (continuous) bilinear map in the two cases (1)  $s < -1/2$  and (2)  $s > 2k - \frac{1}{2}$ .

- If  $s < -1/2$ ,

$$\begin{aligned} \hat{u}_{N,0}(\xi) &= N^{\frac{1}{2}-k} \chi_{[-N-\frac{1}{N}, -N]}(\xi), \\ \hat{n}_{N,0}(\xi) &= N^{\frac{1}{2}-s} \chi_{[2N-1, 2N-1+\frac{1}{N}]}(\xi) + N^{\frac{1}{2}-s} \chi_{[-2N+1-\frac{1}{N}, -2N+1]}(\xi) \end{aligned}$$

and  $n_{N,1} = 0$ , then  $(u_{N,0}, n_{N,0}, n_{N,1})$  is a sequence such that

$$\|(u_{N,0}, n_{N,0}, n_{N,1})\|_X \sim 1,$$

but

$$\|\partial_\gamma^2 u|_{\gamma=0}(t)\|_{H^k} \geq c(t) N^{-s-\frac{1}{2}}.$$

We note that the second term in the definition of  $\hat{n}_{N,0}(\xi)$  is included solely to make  $n_0(x)$  real.

- If  $s > 2k - \frac{1}{2}$  and we set

$$\hat{u}_{N,0}(\xi) = N^{\frac{1}{2}-k} (\chi_{[-N-\frac{1}{N}, -N]}(\xi) + \chi_{[N+1, N+1+\frac{1}{N}]}(\xi))$$

and  $n_{N,0} = 0, n_{N,1} = 0$ , then  $(u_{N,0}, n_{N,0}, n_{N,1})$  is a sequence such that  $\|(u_{N,0}, n_{N,0}, n_{N,1})\|_X \sim 1$  but

$$\|\partial_\gamma^2 n|_{\gamma=0}(t)\|_{H^s} \geq c(t)N^{s-(2k-\frac{1}{2})}$$

The second was considered in §3 as part of the proof of Theorem 1.2, and thus reproduces a weaker version of that result. We now carry out a proof of the first case to establish Theorem 1.4. Since

$$[\partial_\gamma^2 u|_{\gamma=0}(t)](\xi) = \int_0^t e^{-i(t-t')\xi^2} [Uu_{N,0}(t')W(n_{N,0}, 0)(t')](\xi) dt' \tag{6.5}$$

we need to examine

$$\begin{aligned} & [Uu_{N,0}(t')W(n_{N,0}, 0)(t')](\xi) \\ &= \int_{\xi_1} e^{-it'\xi_1^2} \hat{u}_{N,0}(\xi_1) \cos(t'(\xi - \xi_1)) \hat{n}_{N,0}(\xi - \xi_1) d\xi_1 \\ &= (e^{-it'N^2} \cos(t'(2N - 1)) + \mathcal{O}(t')) \int_{\xi_1} \hat{u}_{N,0}(\xi_1) \hat{n}_{N,0}(\xi - \xi_1) d\xi_1 \end{aligned}$$

by the support properties of  $u_{N,0}$  and  $n_{N,0}$ . Directly evaluating the convolution gives

$$(e^{-it'(N-1)^2} + e^{-it'(N^2+2N-1)} + \mathcal{O}(t'))N^{1-k-s}g(\xi)$$

where  $g(\xi) = g_1(\xi) + g_2(\xi)$  consists of two triangular step functions, each of height  $1/N$  and width  $2/N$ , centered at  $N - 1$  and  $-3N + 1$ , respectively. Specifically,

$$g_1(\xi) = \begin{cases} \frac{1}{N} - |\xi - (N - 1)| & \text{if } |\xi - (N - 1)| \leq \frac{1}{N} \\ 0 & \text{otherwise} \end{cases}$$

$$g_2(\xi) = \begin{cases} \frac{1}{N} - |\xi - (-3N + 1)| & \text{if } |\xi - (-3N + 1)| \leq \frac{1}{N} \\ 0 & \text{otherwise} \end{cases}$$

We have by the support properties of  $g_1(\xi)$  and  $g_2(\xi)$  and (6.5),

$$\begin{aligned} & [\partial_\gamma^2 u|_{\gamma=0}(t)](\xi) \\ &= +N^{1-k-s}g_1(\xi) \int_0^t e^{-i(t-t')(N-1)^2} (e^{-it'(N-1)^2} + e^{-it'(N^2+2N-1)}) dt' \\ & \quad + N^{1-k-s}g_2(\xi) \int_0^t e^{-i(t-t')(-3N+1)^2} (e^{-it'(N-1)^2} + e^{-it'(N^2+2N-1)}) dt' \\ & \quad + N^{1-k-s}g(\xi)\mathcal{O}(t^2) \end{aligned}$$

Evaluating each component separately gives

$$[\partial_\gamma^2 u|_{\gamma=0}(t)](\xi) = N^{1-k-s}g_1(\xi)e^{-it(N-1)^2}t + N^{1-k-s}g(\xi)(\mathcal{O}(t^2) + \mathcal{O}(N^{-1}))$$

Thus, provided  $t$  is chosen small and  $N$  sufficiently large, the first term is pointwise dominant, giving

$$\|\partial_\gamma^2 u|_{\gamma=0}(t)\|_{H^k} \geq tN^{-\frac{1}{2}-s}$$

completing the proof.

**Acknowledgments.** I would like to thank Jim Colliander for his clear explanation of how to construct counterexamples to bilinear estimates and for other helpful discussion on this topic. Also, I would like to thank Guixiang Xu for carefully reading §1–3 of the paper and pointing out numerous misprints and an error. Finally, I would like to thank the anonymous referee for providing several helpful suggestions for improvement.

## REFERENCES

- [1] J. Bourgain and J. Colliander, *On wellposedness of the Zakharov system*, Internat. Math. Res. Notices (1996), no. 11, 515–546. MR MR1405972 (97h:35206)
- [2] Björn Birnir, Carlos E. Kenig, Gustavo Ponce, Nils Svanstedt, and Luis Vega, *On the ill-posedness of the IVP for the generalized Korteweg-de Vries and nonlinear Schrödinger equations*, J. London Math. Soc. (2) **53** (1996), no. 3, 551–559. MR1396718 (97d:35233)
- [3] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations*, Geom. Funct. Anal. **3** (1993), no. 2, 107–156. MR1209299 (95d:35160a)
- [4] Michael Christ, James Colliander, and Terence Tao, *Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations*, Amer. J. Math. **125** (2003), no. 6, 1235–1293. MR2018661 (2005d:35223)
- [5] Michael Christ, James Colliander, and Terence Tao, *Ill-posedness for nonlinear Schrödinger and wave equations*, to appear, Ann. Inst. H. Poincaré Anal. Non Linéaire, arxiv.org preprint [math.AP/0311048](https://arxiv.org/abs/math/0311048).
- [6] James Colliander, *The initial value problem for the Zakharov system*, Ph.D. thesis, University of Illinois at Urbana Champaign, 1997.
- [7] J. Ginibre, Y. Tsutsumi, and G. Velo, *On the Cauchy problem for the Zakharov system*, J. Funct. Anal. **151** (1997), no. 2, 384–436. MR1491547 (2000c:35220)
- [8] Bo Ling Guo, *Existence and nonexistence of global solution, soliton solution for a class of the systems of generalized Zakharov equations*, Nonlinear evolutions (Balaruc-les-Bains, 1987), World Sci. Publishing, Teaneck, NJ, 1988, pp. 711–720. MR1031546 (90m:35169)
- [9] Carlos E. Kenig, Gustavo Ponce, and Luis Vega, *On the ill-posedness of some canonical dispersive equations*, Duke Math. J. **106** (2001), no. 3, 617–633. MR1813239 (2002c:35265)
- [10] N. Tzvetkov, *Ill-posedness issues for nonlinear dispersive equations*, arxiv.org preprint [math.AP/0411455](https://arxiv.org/abs/math/0411455).
- [11] Ya Ping Wu, *Orbital stability of solitary waves of Zakharov system*, J. Math. Phys. **35** (1994), no. 5, 2413–2422. MR1271934 (95e:35170)

JUSTIN HOLMER

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840, USA  
E-mail address: [holmer@math.berkeley.edu](mailto:holmer@math.berkeley.edu)