

References:

- Alshamrani, S., & Al-Surimi, K. (2018). Reducing the rate of blood culture contamination in the emergency department of a university teaching hospital. *Global Journal on Quality and Safety in Healthcare*, 1(1), 13–18. https://doi.org/10.4103/jqsh.jqsh_5_18
- Doern, G. V., Carroll, K. C., Diekema, D. J., Garey, K. W., Rupp, M. E., Weinstein, M. P., & Sexton, D. J. (2020). Practical guidance for clinical microbiology laboratories: a comprehensive update on the problem of blood culture contamination and a discussion of methods for addressing the problem. *Clinical Microbiology Reviews*, 33(1). <https://doi.org/10.1128/cmr.00009-19>
- Kai, M., Miyamoto, K., Akamatsu, K., Tsujita, A., & Nishio, M. (2020). Effect of a bundle-approach intervention against blood culture contamination in the emergency department. *Journal of Infection and Chemotherapy*, 26(8), 785–789. <https://doi.org/10.1016/j.jiac.2020.03.005>
- Lippincott, W. (2023). *Lippincott Nursing Procedures*. (9th ed.). Wolters Kluwer.
- Moeller, D. (2017). Eliminating blood culture false positives: Harnessing the power of nursing shared governance. *Journal of Emergency Nursing*, 43(2), 126–132. <https://doi.org/10.1016/j.jen.2016.07.001>
- Skoglund, E., Dempsey, C. J., Chen, H. & Garey, K. W. (2019). Estimated clinical and economic impact through use of a novel blood collection device to reduce blood culture contamination in the emergency department: A Cost-Benefit Analysis. *Journal of Clinical Microbiology*, 57(1). <https://doi.org/10.1128/JCM.01015-18>
- Steed, L. L. (2017, August 20). Reducing blood culture contamination. *Clinical Lab Products*. <https://clpmag.com/disease-states/reducing-blood-culture-contamination/>